



## NEUROBIOLOGY OF ADIPOSE TISSUE

**Gorana Rančić<sup>1</sup>, Mariyana G. Hristova<sup>2</sup>, and Vladmila Bojanic<sup>3</sup>**

<sup>1</sup>Department of Embryology and Histology, Medical Faculty, University of Niš, Niš, Serbia, <sup>2</sup>Division of Cell Biology, Medical University, Varna, Bulgaria, and <sup>3</sup>Department of Pathophysiology, Medical Faculty, University of Niš, Niš, Serbia

*Adipose tissue's secretory phenotype paradigm shift has been upregulating since December 1, 1994, the birthday of leptin, an endocrine signaling protein (adipokine), which triggered the development of adipoendocrinology. Today, more than hundred adipokines are identified. Here an update of adipose-derived neurotrophic factors and neuropeptides and their receptors is presented, raising a hypothesis of neuroendocrine potential of this dynamic tissue. **Biomed Rev 2008; 19: 45-48.***

**Key words:** adipose tissue, neurotrophins, neuropeptides

### INTRODUCTION

Today, adipose tissue (mainly its white phenotype) is increasingly recognized as a dynamic tissue which may, by sending and receiving different types of protein and non-protein signals, communicate with many organs in the human's body. Indeed, an extensive body of work has revealed that adipose tissue expresses not only metabolic, but also endo-, auto-, intra- and paracrine phenotypes. This new biology is achieved predominantly through synthesis, storage, and release (that is, secretion) of adipokines, which include more than hundred highly active signaling proteins abundantly secreted by adipose tissue, particularly when hypertrophied, inflamed or other re-

lated conditions (1-12). These factors may contribute to brain, immune, vascular and metabolic functions, including feeding behavior, memory, learning, circadian rhythm, inflammation, immunity, vascular tone, and insulin resistance.

There is at present evidence that the sharing of ligands (growth factors, cytokines, and adipokines) and their receptors constitute a "global" molecular language of the human's body, including neural, immune and adipose cells. For instance, leptin and adiponectin exert effects on hypothalamus, sympathetic nerves and immune cells (reviewed in 5-7, 13, also see 14-16). Noteworthy, though the brain has been recognized as

Received 22 November 2008, accepted 15 December 2008.

Correspondence and reprint request to Dr Gorana Rančić, Department of Embryology and Histology, Medical Faculty, University of Niš, Niš, Serbia. Tel.: 38 11853 5955, E-mail: goranaranc@medfak.ni.ac.yu

a prominent site of neuropeptide synthesis for more than 30 years, it was recently discovered that adipose-specific genes (e.g. for leptin, resistin and adiponectin) are also expressed in the brain (17,18), and that leptin exerts neuroprotective effect (19,20). Moreover, substance P, neuropeptide tyrosine (NPY), calcitonin gene-related protein, kisspeptin and other neuropeptides (21-26) as well as amino acid neurotransmitters (27) are also produced by adipose tissue. In the same vein, most hypothalamic and pituitary neuropeptides, hormones and releasing factors, termed "adipotrophins" (28), express their receptors in adipose tissue, creating hypothalamic-pituitary-adipose axis (29), and possibly, pineal-adipose axis. Further, adipose stem cells appear to be a promising therapeutic tool in brain injury (30,31). Noteworthy, various neurotrophic factors including nerve growth factor (NGF), brain-derived neurotrophic factor (24,32-36), also vascular endothelial growth factor and angiopoietin (37-39), are produced and released from adipose tissue.

While NGF was first discovered by Rita Levi-Montalcini in 1951 as nerve growth stimulating factor produced in largest amount by submandibular glands (40, also 41 for extraneuronal actions of NGF, and 42,43 for BDNF), it appears today that the adipose tissue might be a major biological source of NGF and other neurotrophic factors as well as neuropeptides and neurotransmitters, selected examples being indicated in Table 1. Altogether, this complex secretory capacity raises a hypothesis of neuroendocrine potential of adipose tissue.

#### ACKNOWLEDGMENTS

The authors appreciate the stimulating discussions with Luigi Aloe and George Chaldakov.

#### REFERENCES

- Zhang Y, Proenca R, Maffei M, Barone M, Leopold L, Friedman JM. Positional cloning of the mouse obese gene and its human homologue. *Nature* 1994; 372: 425-432.
- Chaldakov GN, Stankulov IS, Hristova M, Ghenev PI. Adipobiology of disease: adipokines and adipokine-targeted pharmacology. *Curr Pharm Des* 2003; 9:1023-1031.
- Trayhurn P, Wood IS. Adipokines: inflammation and pleiotropic role of white adipose tissue. *Br J Nutr* 2004; 92, 347-355.
- Yudkin JS, Eringa E, Stehouwer CDA. „Vasocrine“ signalling from perivascular fat: a mechanism linking insulin resistance to vascular disease. *Lancet* 2005; 365: 1817-1820.
- Chaldakov GN, Fantuzzi G, editors. Adipopharmacology of Disease. *Biomed Rev* 2006; 17: 1-128.
- Chaldakov GN, editor. The Adipobiology of Disease. *Immun Endocr Metab Agents Med Chem* 2007; 7: 105-173.
- Fantuzzi G, Mazzone T, editors. *Nutrition and Health: Adipose Tissue and Adipokines in Health and Disease*. Humana Press Inc., Tokowa, NJ. 2007.
- Iacobellis G, Sharma AM. Epicardial adipose tissue as new cardio-metabolic risk marker and potential therapeutic target in the metabolic syndrome. *Curr Pharm Des* 2007; 13: 2180-2184.
- Sacks H, Fain JN. Human epicardial adipose tissue: A review. *Am Heart J* 2007; 153: 907-917.

**Table 1.** Adipose tissue-produced factors affecting nerve cells\*

#### Neuropeptides

Neuropeptide tyrosine (NPY)  
Substance P  
Calcitonin gene-related peptide  
Adrenomedullin  
Somatostatin  
Agouti protein  
Kisspeptin

#### Neurotrophic factors

Nerve growth factor  
Brain-derived neurotrophic factor  
Ciliary neurotrophic factor  
Metallothioneins  
Glial cell line-derived neurotrophic factor  
Angiopoietin-1  
Vascular endothelial growth factor

#### Neurotransmitters

Glutamate  
Gamma-aminobutyric acid (GABA)

#### Others

Leptin  
Adiponectin  
Insulin-like growth factor  
Free fatty acids

\*References included 14-16, 21-29, 32-38, 44.

10. Guzik TJ, Marvar PJ, Czesnikiewicz-Guzik M, Korb R. Perivascular adipose tissue as a messenger of the brain-vascular axis: role in vascular inflammation and dysfunction. *J Physiol Pharmacol* 2007; 58: 591-610.
11. Wang P, Mariman E, Renes J, Keijer J. The secretory function of adipocytes in the physiology of white adipose tissue. *J Cell Physiol* 2008; 216: 3-13.
12. Wang B, Wood IS, Trayhurn P. PCR arrays identify metallothionein-3 as highly hypoxia-inducible gene in human adipocytes. *Biochem Biophys Res Commun* 2008; 368: 88-93.
13. Pond CM. Adipose tissue and the immune system. *Prostaglandins Leukot Essent Fatty Acids* 2005; 73: 17-30.
14. Guerre-Millo M. Adiponectin: an update. *Diabetes Metab* 2008; 34: 12-18.
15. Tanida M, Shen J, Horii Y, Matsuda M, Kihara S, Funahashi T, et al. Effects of adiponectin on the renal sympathetic nerve activity and blood pressure in rats. *Exp Biol Med (Maywood)* 2007; 232: 390-397.
16. Coope A, Milanski M, Araújo EP, Tambascia M, Saad MJ, Geloneze B, et al. AdipoR1 mediates the anorexigenic and insulin/leptin-like actions of adiponectin in the hypothalamus. *FEBS Lett* 2008; 582: 1471-1476.
17. Wilkinson M, Brown R, Imran SA, Ur E. Adipokine gene expression in brain and pituitary gland. *Neuroendocrinology* 2007; 86: 191-209.
18. Brown R, Thompson HJ, Imran SA, Ur E, Wilkinson M. Traumatic brain injury induces adipokine gene expression in rat brain. *Neurosci Lett* 2008; 432: 73-78.
19. Zhang F, Chen J. Leptin protects hippocampal CA1 neurons against ischemic injury. *J Neurochem* 2008; 107: 578-587.
20. Tang BL. Leptin as a neuroprotective agent. *Biochem Biophys Res Commun* 2008; 368:181-185.
21. Kos K, Harte AL, James S, Snead DR, O'Hare JP, McTernan PG, et al. Secretion of neuropeptide Y in human adipose tissue and its role in maintenance of adipose tissue mass. *Physiol Endocrinol Metab* 2007; 293: E1335-1340.
22. Gupta P, Harte AL, da Silva NF, Khan H, Barnett AH, Kumar S, et al. Expression of calcitonin gene-related peptide, adrenomedullin, and receptor modifying proteins in human adipose tissue and alteration in their expression with menopause status. *Menopause* 2007; 14: 1031-1038.
23. Seboek D, Linscheid P, Zulewski H, Langer I, Christ-Crain M, Keller U, et al. Somatostatin is expressed and secreted by human adipose tissue upon infection and inflammation. *J Clin Endocrinol Metab* 2004; 89: 4833-4839.
24. Hausman GJ, Barb CR, Dean RG. Patterns of gene expression in pig adipose tissue: Insulin-like growth factor system proteins, neuropeptide Y (NPY), NPY receptors, neurotrophic factors and other secreted factors. *Domest Anim Endocrinol* 2008 February 12; In press.
25. Voisey J, van Daal A. Agouti: from mouse to man, from skin to fat. *Pigment Cell Res* 2002; 15: 10-18.
26. Brown RE, Imran SA, Ur E, Wilkinson M. KiSS-1 mRNA in adipose tissue is regulated by sex hormones and food intake. *Mol Cell Endocrinol* 2008; 281:64-72.
27. Nicolaysen A, Gammelsaeter R, Storm-Mathisen J, Gundersen V, Iversen PO. The components required for amino acid neurotransmitter signaling are present in adipose tissue. *J Lipid Res* 2007; 48: 2123-2132.
28. Schäffler A, Schölmerich J, Buechler C. The role of "adipotrophins" and the clinical importance of a potential hypothalamic-pituitary-adipose axis. *Nat Clin Pract Endocrinol Metab* 2006; 2: 374-383.
29. Schäffler A, Schölmerich J, Buechler C. Hypothesis paper. Brain talks with fat – evidence for a hypothalamic-pituitary-adipose axis? *Neuropeptides* 2005; 39: 363-367.
30. Ohta Y, Takenaga M, Tokura Y, Hamaguchi A, Matsumoto T, Kano K, et al. Mature adipocyte-derived cells, dedifferentiated fat cells (DFAT), promoted functional recovery from spinal cord injury-induced motor dysfunction in rats. *Cell Transplant* 2008; 17: 877-886.
31. Wei X, Du Z, Zhao L, Feng D, Wei G, He Y, et al. IFATS Series: The conditioned media of adipose stromal cells protect against hypoxia-ischemia-induced brain damage in neonatal rats. *Stem Cells* 2008 Nov 20. In press.
32. Chaldakov GN, Fiore M, Stankulov IS, Manni L, Hristova MG, Antonelli A, et al. Neurotrophin presence in human coronary atherosclerosis and metabolic syndrome: a role for NGF and BDNF in cardiovascular disease? *Prog Brain Res* 2004; 146:279-289.
33. Wang B, Jenkins JR, Trayhurn P. Expression and secretion of inflammation-related adipokines by human adipocytes differentiated in culture: integrated response to TNF- $\alpha$ . *Am J Physiol Endocrinol Metab* 2005; 288: E731-E740.
34. Bulló M, Peeraully MR, Trayhurn P. Stimulation of NGF expression and secretion in 3T3-L1 adipocytes by prostaglandins PGD2, PGJ2, and Delta12-PGJ2. *Am J Physiol Endocrinol Metab* 2005; 289, E62-67.
35. Sornelli F, Fiore M, Chaldakov GN, Aloe L. Brain-

- dervied neurotrophic factor: a new adipokine. *Biomed Rev* 2007; 18: 85-88.
36. Ryan VH, German AJ, Wood IS, Hunter L, Morris P, Trayhurn P. NGF gene expression and secretion by canine adipocytes in primary culture: Upregulation by the inflammatory mediators LPS and TNFalpha. *Horm Metab Res* 2008; 40: 861-868.
  37. Hansen-Algenstaedt N, Algenstaedt P, Schaefer C, Hamann A, Wolfram L, Cingöz G, *et al.* Neural driven angiogenesis by overexpression of nerve growth factor. *Histochem Cell Biol* 2006; 125: 637-649.
  38. Kosacka J, Nowicki M, Kacza J, Borlak J, Engele J, Spanel-Borowski K. Adipocyte-derived angiopoietin-1 supports neurite outgrowth and synaptogenesis of sensory neurons. *J Neurosci Res* 2006; 83: 1160-1169.
  39. Lazarovici P, Marcinkiewicz C, Lelkes PI. Cross talk between the cardiovascular and nervous systems: neurotrophic effects of vascular endothelial growth factor (VEGF) and angiogenic effects of nerve growth factor (NGF) – implications in drug development. *Curr Pharm Des* 2006; 12: 2609-2622.
  40. Levi-Montalcini R. The nerve growth factors 35 years later. *Science* 1987; 237: 1154-1162.
  41. Aloe L, Tirassa P, Bracci-Laudiero L. Nerve growth factor in neurological and non-neurological diseases: basic findings and emerging pharmacological prospectives. *Curr Pharm Des* 2001; 7: 113-123.
  42. Tsukinoki K, Saruta J, Sasaguri K, Miyoshi Y, Jinbu Y, Kusama M, *et al.* Immobilization stress induces BDNF in rat submandibular glands. *J Dent Res* 2006; 85, 844-848.
  43. Tsukinoki K, Saruta J, Muto N, Sasaguri K, Sato S, Tan-Ishii N, *et al.* Submandibular glands contribute to increases in plasma BDNF levels. *J Dent Res* 2007; 86, 260-264.
  44. Ma D, Lu L, Boneva NB, Warashina S, Kaplamadzhiev DB, Mori Y, *et al.* Expression of free fatty acid receptor GPR40 in the neurogenic niche of adult monkey hippocampus. *Hippocampus* 2008;18:326-333.